



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Put $m=1$; then a =any value between $a>3$ and $a=10\frac{1}{2}$.

Put $a=4$; then $n=4$, $p=7$,

$x=13\frac{2}{3}$ and $y=\frac{1}{3}$;

$\therefore mx=13\frac{2}{3}$, $nz=54\frac{2}{3}$, and $px=95\frac{2}{3}$.

Put $m=2$; then a =any value between $a>3$ and $a=5\frac{1}{2}$.

Put $a=4$; then $n=8$, $p=14$,

$x=7$ and $y=\frac{1}{2}$;

$\therefore mx=14$, $nz=56$, and $px=98$.

For want of space, we will let the reader solve for sets of values when $m=0$, $m>4$, and $a<3$. The discussion of these values may prove interesting.

Also solved by *H. W. DRAUGHON*.

13. Proposed by *H. C. WHITAKER, B. S., M. E., Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.*

Six city boys, Jim, Josh, Jerry, Jack, Jake and Jearje went into the country to steal apples from a tree. While three kept watch, the other three climbed up and got what they wanted. Then they came down while the other three rascals went up and stole. The one that got most was one of the last to go up.

Each trio of thieves took the same number and had each boy taken as many as he did take in *each of that number of pockets*, each trio would also have taken the same number and the tree would have lost 538 apples. As it was, Josh got more than Jack, but Jearje got as many as Josh and Jack together, while Jake got twice as many as Jerry and two more than Jim. What were the names of the three that first kept watch?

Solution by *H. W. DRAUGHON, Clinton, Louisiana.*

Let x =Jack's share, y =Josh's share and z =Jerry's share. Then from the conditions, $x+y$ =Jearje's share, $2z$ =Jake's share, and $2z-2$ =Jim's share. To find each share in integers, we must separate $\frac{1}{2} \times 538 = 269$,—the numbers each trio would have taken, had the number in each share been squared, into 2 sets of 3 square numbers the sum of the roots in the two sets being equal. We easily find the required numbers to be, $(13)^2$, $(8)^2$, $(6)^2$, and $(12)^2$, $(10)^2$, $(5)^2$. Since the greatest root is 13, the boys who took respectively 13, 8, and 6 apples compose the trio who first kept watch. Comparing our results with the expressions for each boy's share given above, we find,

First trio;—Jerry, Jearje, and Josh; respective shares;—6, 13, 8. Second trio;—Jack, Jake, and Jim; respective shares;—5, 12, and 10.

\therefore Jerry, Josh and Jearje first kept watch.

Also solved by *A. L. FOOTE, H. C. WHITAKER, and G. B. M. ZERR.*

PROBLEMS.

21. Proposed by *Professor J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.*

A tobacconist has two kinds of smoking tobacco, of which the price of the